

Cluster FGM daily calibration

Instrument

Parameters

Effects

Procedure

Variation

D. Constantinescu¹ Evelyn Liebert² K-H. Fornacon²

¹Institute for Space Sciences, Bucharest

²Institute for Geophysics and Extraterrestrial Physics, Braunschweig

- 1 The Cluster FGM instrument
- 2 Calibration Parameters
- 3 Effects on power spectra
- 4 The daily calibration procedure
- 5 Long term variation

The Cluster FGM instrument(s)

Instrument

Parameters

Effects


Procedure

Variation

- (inboard + outboard) $\times 4$
- very stable during the (long) mission
- however, daily calibration is necessary
- uses ranges depending on field magnitude

| range | B_{\max} (nT) | resolution (nT) | |
|-------|-----------------|-----------------|------------|
| 2 | 64 | 1/128 | |
| 3 | 256 | 1/32 | |
| 4 | 1024 | 1/8 | |
| 5 | 4096 | 1/2 | since 2006 |
| 6 | 16384 | 2 | since 2008 |
| 7 | 65536 | 8 | since 2009 |

Calibration parameters



12 independent parameters

- 3 elevation angles θ_i
- 3 azimuth angles φ_i
- 3 gains G_i
- 3 offsets O_i

$$\begin{bmatrix} B'_x \\ B'_y \\ B'_z \end{bmatrix} = \mathcal{M} \cdot \begin{bmatrix} B_x \\ B_y \\ B_z \end{bmatrix} + \begin{bmatrix} O_x \\ O_y \\ O_z \end{bmatrix}$$

$$\mathcal{M} = \begin{bmatrix} G_x \sin \theta_x \cos \varphi_x & G_x \sin \theta_x \sin \varphi_x & G_x \cos \theta_x \\ G_y \sin \theta_y \cos \varphi_y & G_y \sin \theta_y \sin \varphi_y & G_y \cos \theta_y \\ G_z \sin \theta_z \cos \varphi_z & G_z \sin \theta_z \sin \varphi_z & G_z \cos \theta_z \end{bmatrix}$$

Small orthogonality deviations

Change variables

$$\theta_x \rightarrow \theta_x \quad \theta_y \rightarrow 90 - \theta_y \quad \theta_z \rightarrow 90 - \theta_z$$

$$\varphi_x \rightarrow \varphi_x \quad \varphi_y \rightarrow \varphi_y \quad \varphi_z \rightarrow 90 + \varphi_z$$

Take differences

$$\Delta G_{yz} = G_y - G_z \quad \text{spin plane gain mismatch}$$

$$\Delta \varphi_{yz} = \varphi_y - \varphi_z \quad \text{spin plane nonorthogonality}$$

Calibration matrix in the first order:

$$\mathcal{M} = \begin{bmatrix} G_x & G_x \theta_x \sin \varphi_x & G_x \theta_x \cos \varphi_x \\ G_y \theta_y & G_y & -G_y \varphi_y \\ G_y \theta_z & G_y (\varphi_y + \Delta \varphi_{yz}) & G_y + \Delta G_{yz} \end{bmatrix}$$

Finding the right parameters

1 Fourier power spectrum [Kepko et al 1996]

- errors lead to coherent signals:

- spin plane $\theta_y, \theta_z, O_y, O_z$ $\rightarrow \omega_{\text{spin}}$
- spin plane $\Delta\varphi_{yz}, \Delta G_{yz}$ $\rightarrow 2\omega_{\text{spin}}$
- spin axis θ_x, φ_x $\rightarrow \omega_{\text{spin}}$


2 Solar wind calibration [Hedgecock 1975]

- assume no correlation between \mathbf{B} and \mathbf{e}_B
 - spin axis O_x

3 Range change continuity

- both \mathbf{B} and its time derivative
-


Error effects: spin tone (yz)



- spin plane elevation angles
- spin plane offsets

$$\begin{bmatrix} G_x & G_x \theta_x \sin \varphi_x & G_x \theta_x \cos \varphi_x \\ G_y \theta_y & G_y & -G_y \varphi_y \\ G_y \theta_z & G_y (\varphi_y + \Delta \varphi_{yz}) & G_y + \Delta G_{yz} \end{bmatrix} \cdot \begin{bmatrix} B_x \\ B_y \\ B_z \end{bmatrix} + \begin{bmatrix} O_x \\ O_y \\ O_z \end{bmatrix}$$


Error effects: spin tone (x)



- spin axis alignment
- the angles are coupled

$$\begin{bmatrix} G_x & G_x \theta_x \sin \varphi_x & G_x \theta_x \cos \varphi_x \\ G_y \theta_y & G_y & -G_y \varphi_y \\ G_y \theta_z & G_y (\varphi_y + \Delta \varphi_{yz}) & G_y + \Delta G_{yz} \end{bmatrix} \cdot \begin{bmatrix} B_x \\ B_y \\ B_z \end{bmatrix} + \begin{bmatrix} O_x \\ O_y \\ O_z \end{bmatrix}$$

Error effects: 2nd harmonic



- spin plane nonorthogonality
- spin plane gain mismatch

$$\begin{bmatrix} G_x & G_x \theta_x \sin \varphi_x & G_x \theta_x \cos \varphi_x \\ G_y \theta_y & G_y & -G_y \varphi_y \\ G_y \theta_z & G_y (\varphi_y + \Delta \varphi_{yz}) & G_y + \Delta G_{yz} \end{bmatrix} \cdot \begin{bmatrix} B_x \\ B_y \\ B_z \end{bmatrix} + \begin{bmatrix} O_x \\ O_y \\ O_z \end{bmatrix}$$

The remaining parameters


The spin axis offset is determined from SW calibration

$$\begin{bmatrix} G_x & G_x \theta_x \sin \varphi_x & G_x \theta_x \cos \varphi_x \\ G_y \theta_y & G_y & -G_y \varphi_y \\ G_y \theta_z & G_y (\varphi_y + \Delta \varphi_{yz}) & G_y + \Delta G_{yz} \end{bmatrix} \cdot \begin{bmatrix} B_x \\ B_y \\ B_z \end{bmatrix} + \begin{bmatrix} O_x \\ O_y \\ O_z \end{bmatrix}$$

still undetermined:


- spin axis gain G_x
- one of the spin plane gains G_y
- spin phase difference φ_y

Daily calibration procedure




- mix of
 - C
 - Fortran
 - IDL
 - perl
- one month
- SP off: automatic
- other: manual

Long term variation: C1 SP off




offset drift: O_y : 0.2 nT/yr; O_z : 0.1 nT/yr
seasonal variation: temperature related?

Long term variation: C234 SP off







very little (if any) offset drift
visible seasonal variation

Long term variation: C1r3 off




Long term variation: C1r4567 off



Long term variation: C1 \perp & gain

Instrument
Parameters
Effects
Procedure
Variation



Summary

- 12 independent calibration parameters
- 8 calibration parameters from Fourier spectra
- 1 calibration parameter from the SW cal
- adjustment for rc continuity
- 0.1 - 0.2 nT/yr offset drift for C1
- much smaller drift for the other sc
- seasonal variation of offsets for all sc

Acknowledgement

This study is supported by the TUNED project financed by the STAR program of the Romanian Space Agency